

Walter K. Nahm, M.D., Ph.D.

Diplomate of the American Board of Dermatology
American College of Mohs Micrographic Surgery
And Cutaneous Oncology
American Academy of Cosmetic Surgery

Dermatology
Mohs Micrographic Surgery
Laser Surgery
Cosmetic Surgery

INSTRUCTIONS FOR THE AREA WHERE THE GRAFT WAS TAKEN FROM

Donor site location: _____

Donor site sutured

1. Bulky pressure dressing to remain intact for 24 hours
2. After 24 hours remove the bulky bandages. Leave the flat dressing on your skin until the dressing is removed approximately 7 days following surgery.
3. Original piece of tape to remain intact until the sutures are removed. If the tape becomes loose, reinforce it with another piece of tape,

Donor site, naturally healing wound

1. Bulky dressing to remain intact for 24 hours
2. After 24 hours remove the dressing and clean with soapy water, using a gauze pad.....soak for approximately 5 minutes.
3. Dry with a sterile gauze pad. **Apply Polysporin, Bacitracin, Aquaphor, or Vaseline** (all of which can be purchased over the counter) over the entire wound, cover the wound with a telfa dressing pad and micropore tape (paper tape) Continue daily for 2 - 3 weeks or until the wound heals.

IN CASE OF AN EMERGENCY:

During regular office hours call (858) 278-8835.....Walter K. Nahm, M.D., Ph.D

After hours and on weekends, you can call the doctor on his cell phone, (619) 318-0952, if no answer, call (858) 278-8835.

Specializing in Cosmetic & Medical Dermatology

7695 Cardinal Court, Suite 200 • San Diego, CA 92123 • Phone: 858.278.8835 • Fax: 858.386.4776